arXiv:1212.3195v1 [q-fin.ST] 13 Dec 2012

Non stationary multifractality in stock returns

Raffaello Morales and T. Di Matteo
Department of Mathematics, King's College London, Strand, WC2R 2LS, London, UK

Tomaso Aste*
Department of Computer Science, University College London, Gower Street, WC1E 6BT, London, UK

We report evidence that empirical data show time varying multifractal properties. This is obtained by comparing empirical observations of the weighted generalised Hurst exponent (wGHE) with time series simulated via Multifractal Random Walk (MRW) by Bacry *et al.* [*E.Bacry, J.Delour and J.Muzy, Phys.Rev.E* **64** *026103, 2001*]. While dynamical wGHE computed on synthetic MRW series is consistent with a scenario where multifractality is constant over time, fluctuations in the dynamical wGHE observed in empirical data fail to be in agreement with a MRW with constant intermittency parameter. This is a strong argument to claim that observed variations of multifractality in financial time series are to be ascribed to a structural breakdown in the temporal covariance structure of stock returns series. As a consequence, multi fractal models with a constant intermittency parameter may not always be satisfactory in reproducing financial market behaviour.

Keywords: Multifractality, Generalized Hurst exponent, Multifractal models.

I. INTRODUCTION

The concept of multifractality in the context of finance has received much attention in the econophysics literature particularly over the last two decades. Many empirical studies have investigated financial data scaling behaviour [1–7] and several models have been proposed to account for the observed multifractal features [8–15]. Multifractal behaviour has consequently become a downright stylised fact of financial market data [16], being observed across several classes of assets: from daily stock returns to foreign exchange rates and composite indices [2, 5, 17–19]. If one denotes p_t the asset price at time t and $r_{t,\tau} = p_{t+\tau} - p_t$ the return at time t and scale τ, then the process $r_{t,\tau}$ is multifractal if the following scaling is observed [5]

$$\mathbb{E}(|r_{t,\tau}|^q) = c_q \tau^{\zeta_q}, \tag{1}$$

where c_q is a constant and ζ_q is a non-linear function, called the scaling function. The departure from linearity of ζ_q is what distinguishes multifractal processes from uni-scaling ones.

Multifractality is also particularly appealing for modelling financial markets as it offers a simple behavioural interpretation: looking at the volatility at different time scales is a very natural way to assess the impact of heterogeneous agents in the market and therefore any measure of scaling behaviour can convey information about the efficiency of a given market [20]. In this regard some authors have suggested and confirmed through extensive empirical studies, that scaling exponents can be representative of the stage of development of a market [2, 17]. The same works have convincingly shown that, through a hierarchy of scaling exponents, it is possible to classify markets according to their degree of development: emerging (developed) markets exhibit scaling exponents significantly larger (smaller) that those expected for purely uncorrelated noise.

In the aforementioned spirit, the time evolution of the scaling function can be useful to track time varying properties of the market. For this reason ζ_q has also been studied dynamically via the time dependent Hurst exponent [4, 21, 22]. In a recent publication [23] the authors have observed large fluctuations in multifractality measured via the generalised Hurst exponent in empirical daily data [23] across different stock sectors. When one is faced with the task of ascertaining the nature of dynamical fluctuations in these quantities, the subtle issue is being able to distinguish between bare statistical fluctuations, which are due to the finiteness of the sample, and true structural changes in the underlying multifractal process. In this paper we study the problem of validating dynamical fluctuations of the scaling functions, performing an extensive analysis of of empirical stock returns and comparing their properties with those of synthetic multifractal series. Among all existing models we focus on the Multifractal Random Walk (MRW) introduced by Bacry *et al.* [8] because of its parsimonious formulation and its success in the econophysics literature.

The paper is organised as follows: in Section II, after giving a short summary of the main tools used in this context, namely the dynamical GHE method and an overview of the MRW model, we illustrate the relationship between the scaling exponents and the spectrum of the model. In Section III we report the main findings on the varying multifractality of empirical stock returns data, which are backed up by a simulation study on synthetic MRW series in Section IV; a summary and conclusive remarks are drawn in Section V.

---

* t.aste@ucl.ac.uk

II. GENERALIZED HURST EXPONENT AND MULTIFRACTAL RANDOM WALK

The MRW model [8] is constructed, in its discrete version, via the sum of stationary increments

$$X_{\Delta t}(t) = \sum_{k=1}^{t/\Delta t} x_{\Delta t,k}, \tag{2}$$

where each contribution is defined in the following way:

$$x_{\Delta t,k} = \epsilon_{\Delta t,k}\, e^{\omega_{\Delta t,k}}. \tag{3}$$

A continuous multifractal model is obtained in the limit $\Delta t \to 0$, but we will only consider the discrete case. In equation (3) $\epsilon_{\Delta t,k}$ is a Gaussian white noise with variance $\sigma \cdot \Delta t$ and $e^{\omega_{\Delta t,k}}$ represents the stochastic volatility, uncorrelated with ϵ. By taking $\omega_{\Delta t,k}$ as a stationary gaussian process, we have log-normal volatility components. What distinguishes the limit $\Delta t \to 0$ of $X_{\Delta t}(t)$ from a Brownian motion is the choice of the correlation structure of the process $\omega_{\Delta t,k}$, which is chosen, according to cascade-like processes [24], as

$$\mathrm{Cov}(\omega_k, \omega_{k+h}) = \lambda^2 \log \frac{T}{(1+|h|)\Delta t} \tag{4}$$

as long as it remains positive, and zero otherwise. The logarithmic decay with lag h of the auto covariance creates long memory in the process. This specification implicitly defines the correlation length T and the intermittency parameter λ. The appeal of this model for describing stock returns evolution lies in its ability to reproduce faithfully the most common stylised facts of financial markets. Its stationarity and its causal structure make it also preferable to other multifractal models [25]. Among the three parameters of the model λ, T and σ, λ is the one governing the non-linearity of the ζ_q spectrum of MRW via the relation [8]:

$$\zeta_q = (q - q(q-2)\lambda^2)/2. \tag{5}$$

The following relation between the generalised Hurst exponent (see Appendix A for details) and the spectrum of ζ_q defined in equation (5) can be established

$$\zeta_q = qH(q). \tag{6}$$

Since $H(q)$ depends only on λ in equation (6), any time dependence for $H(q)$ should be reflected in a time dependence for λ. By considering $q = 1, 2, 3, 4$, equations (5) and (6) the following expressions

$$\begin{aligned} H(1) &= \frac{1}{2}(1+\lambda^2), \qquad H(2) = \frac{1}{2}, \\ H(3) &= \frac{1}{2}(1-\lambda^2), \qquad H(4) = \frac{1}{2}(1+2\lambda^2). \end{aligned} \tag{7}$$

Values of these expressions for different λ's are reported in table I. We perform the following Monte Carlo study in order to compare theoretical predictions with GHE's computed on synthetic time series: for a given value of λ, we simulate 5000 MRW series and we compute the corresponding GHE's for every synthetic series; then, assuming these GHE's are i.i.d., which is quite reasonable since every MRW simulation is independent from the others, we obtain a sample distribution for the GHE's. From this distribution we compute the $\{2.5\%, 50\%, 97.5\%\}$-quantiles which give the range of fluctuation of the GHE beyond which any observation can be deemed anomalous. We perform this study for $q = \{1, 2, 3, 4\}$. Results of this analysis for three different values of λ are reported in table II. By comparing these values with those expected from the MRW model in table I we can conclude that all values computed via the equations given in (7) are compatible with the statistical confidence intervals computed on the synthetic time series. Nonetheless, one can see that confidence intervals for GHE are quite large and do not impose strict bounds to the values retrieved from the model (see a further discussion in Appendix B). The quantiles for $H(1)$ and $H(2)$ reveal no relevant dependence on λ, while a slight dependence is spotted on the 2.5%-quantiles of $H(3)$ and $H(4)$, with the quantile decreasing along with an increase of λ.

TABLE I. Values of $H(q)$ for $q = \{1, 2, 3, 4\}$ computed through the expressions in (7).

	$\lambda = 0.2$	$\lambda = 0.3$	$\lambda = 0.35$
H(1)	0.52	0.5450	0.5613
H(2)	0.50	0.50	0.50
H(3)	0.48	0.4550	0.4387
H(4)	0.46	0.41	0.3775

III. ANALYSIS OF EMPIRICAL DATA

We now turn to show the main findings of the present paper, namely that stock returns exhibit time varying multifractal properties. We analysed daily stock data quoted in the NYSE in the period ranging from 01-01-1995 to 22-10-2012. The data were provided by Bloomberg. In [23] we argued that observed variations in GHE over time could be associated to a transition between different regimes of stability for the firms studied. We have also considered as a measure of multifractal behaviour the quantity

$$\Delta H^w(q, q^{'}) = H^w(q) - H^w(q^{'}), \qquad q \neq q^{'}, \tag{8}$$

where $H^w(q)$ is the weighted Generalized Hurst Exponent (wGHE) [23]. It was pointed out that the dynamical evolution of $\Delta H^w(q, q^{'})$ (which we label $\Delta H^w_t(q, q^{'})$) is useful to track possible changes in multifractality occurring over time. To test the significance of this statement with respect to a multifractal model for stock returns, we

TABLE II. We report numerical results of the quantiles obtained from the distributions of the GHE's computed on several synthetic MRW time series for $H(q)$, $q = 1, 2, 3, 4$. While $H(1)$ and $H(2)$ look quite stable across different values of λ, one may note a dependence on λ particularly in the 2.5%-quantile of $H(3)$ and $H(4)$.

		$\{2.5\%, 50\%, 97.5\%\}$quantiles			
		H(1)	H(2)	H(3)	H(4)
$\lambda = 0.2$	T=2500	$\{0.4256, 0.5046, 0.5870\}$	$\{0.4260, 0.4998, 0.5794\}$	$\{0.4127, 0.4939, 0.5799\}$	$\{0.4050, 0.4840, 0.5852\}$
	T=5000	$\{0.4293, 0.5023, 0.5749\}$	$\{0.4197, 0.4970, 0.5736\}$	$\{0.4107, 0.4924, 0.5726\}$	$\{0.4039, 0.4831, 0.5762\}$
$\lambda = 0.3$	T=2500	$\{0.4281, 0.5105, 0.5911\}$	$\{0.4152, 0.4995, 0.5867\}$	$\{0.3936, 0.4830, 0.5803\}$	$\{0.3831, 0.4753, 0.5891\}$
	T=5000	$\{0.4266, 0.5081, 0.5839\}$	$\{0.4241, 0.4981, 0.5892\}$	$\{0.4017, 0.4861, 0.5806\}$	$\{0.3778, 0.4734, 0.5722\}$
$\lambda = 0.35$	T=2500	$\{0.4271, 0.5112, 0.5975\}$	$\{0.4167, 0.4984, 0.5918\}$	$\{0.3891, 0.4859, 0.5935\}$	$\{0.3653, 0.4747, 0.5832\}$
	T=5000	$\{0.4212, 0.4986, 0.5830\}$	$\{0.4280, 0.5113, 0.5856\}$	$\{0.3899, 0.4814, 0.5801\}$	$\{0.3594, 0.4666, 0.5684\}$



FIG. 1. *The scaling function for Microsoft Corp daily data is compared to that of a notorious unifractal model, i.e. the fractional Brownian motion. The choice of $H = 0.75$ means we are considering a process with long memory although unifractal.*

proceed as follows: first of all we verify that the scaling functions of the data are indeed concave (see figure 1 for an example, where we report the scaling function of the daily stock returns of Microsoft Corp compared to that of a fractional Brownian motion). Then, after estimating the parameters of the model from empirical data, we generate a large number of synthetic MRW time series and for each sample compute the GHE's for $q = 1, 2$, from which we get $\Delta H^w(1,2)$. This endows us with a sample distribution of independent $\Delta H^w(1,2)$'s. From the cumulative distribution function of $\Delta H^w(1,2)$'s one can extract confidence intervals. The latters provide the range of variation beyond which observed fluctuations can be considered to be significant with respect to the underlying model.

The parameters of the model can be extracted from the empirical data through the behaviour of the volatility autocorrelation function. Consider $r_{t,0} = r_t$, where $r_{t,\tau}$ was defined in the introduction, and define

$$y_t = \log|r_t| = \log\sigma_t + \log|\epsilon_t|, \tag{9}$$

Then

$$y_t - y_{t+l} = \log\left(\frac{\sigma_t}{\sigma_{t+l}}\right) + \log|\epsilon_t| - \log|\epsilon_{t+l}|. \tag{10}$$

We can thus compute

$$\begin{aligned}\left\langle (y_t - y_{t+l})^2 \right\rangle &= \langle \left(\log\frac{\sigma_t}{\sigma_{t+l}}\right)^2 \rangle + \langle (\log|\epsilon_t| - \log|\epsilon_{t+l}|)^2 \rangle \\ &= C(l) + A(1 - \delta_{l,0}).\end{aligned} \tag{11}$$

Therefore, $\left\langle (y_t - y_{t+l})^2 \right\rangle$ is proportional, up to a shift given by a constant A, to the log-volatility correlation function. In the MRW model we have for the log-volatility

$$C(l) \simeq -\lambda^2 \log l + A\left(\frac{T}{\Delta t}\right)' \tag{12}$$

and thus we can study $\left\langle (y_t - y_{t+l})^2 \right\rangle$ vs $\log l$ to get an estimate of λ^2 as well as of the correlation length T. A' is another constant which depends on the ratio $T/\Delta t$ and therefore is used to estimate T. The other parameter of the MRW, i.e. σ, can be computed from $\mathrm{Var}(r_t) = \sigma$. We thus estimate both T and λ via linear fits of the $\left\langle (y_t - y_{t+l})^2 \right\rangle$ as function of $\log l$. Values of the parameters obtained for different stocks are reported in table III, together with the quantiles obtained from the empirical distributions of the GHE's. One can appreciate that the quantiles do not seem to vary much across different stocks. In table III we also report the percentage of $\Delta H^w_t(1,2)$ exceeding the quantiles. The rather high number of points falling outside the confidence intervals confirm unassailably that empirical data systematically refuse the hypothesis of constant multifractality. In figure 2 we show the results of this study for several daily stock prices, where $\Delta H^w_t(1,2)$ is shown to exceed the quantiles many times. Note that both IBM and Microsoft

TABLE III. *The parameters of the MRW obtained for a number of stocks from our database, computed by means of the method described in the text. Note the very large values obtained for T, which are in principle difficult to justify given that the length of the time series is 3360 days. Neveretheless we have verified that shorter correlation lengths (up to the order of 10^2) do not make the extreme quantiles larger and hence do not jeopardise the validity of the results.*

	λ	T (days)	σ	Quantiles	Exceedances percentage
Boeing Corp	0.1455	608	1.0651	$\{-0.0205, 0.0039, 0.0265\}$	30 %
Microsoft Corp	0.2436	739	0.6064	$\{-0.0210, 0.0072, 0.0316\}$	53 %
PNC Financials	0.2839	538	1.903	$\{-0.0206, 0.0098, 0.0349\}$	15 %
Sara Lee Corp	0.222	746	0.2937	$\{-0.0204, 0.0071, 0.0318\}$	47.7%
Rowan Cos INC	0.2046	1080	0.7992	$\{-0.0215, 0.0066, 0.0296\}$	11.4%
IBM Corp	0.2363	816	1.7811	$\{-0.0214, 0.0068, 0.0310\}$	39%
Wells Fargo	0.3490	631	0.5923	$\{-0.0237, 0.0134, 0.0433\}$	9%
American Express	0.3278	520	0.8356	$\{-0.0212, 0.0125, 0.0429\}$	14%
General Motors Corp	0.1727	440	1.0291	$\{-0.0211, 0.0063, 0.0278\}$	41%
Citigroup	0.3798	568	6.4585	$\{-0.0202, 0.0146, 0.0458\}$	13%
JPMorgan Chase	0.2950	534	0.9792	$\{-0.0219, 0.0097, 0.0356\}$	27 %
Bank of America	0.3445	670	0.65	$\{-0.0256, 0.0121, 0.0420\}$	30 %
Morgan Stanley	0.2528	538	1.1185	$\{-0.0221, 0.0079, 0.0333\}$	14 %

show a sudden decrease of multifractality corresponding to the 2007-2008 financial crisis, in apparent anticorrelation with JP Morgan's $\Delta H_t^w(1,2)$. This finding is in agreement with what found in [23], where we observed that stocks from the financial sector exhibited increasing multifractality corresponding to the credit crisis.

Empirical quantiles reported in table III and in figure and 2 have been computed from a Monte Carlo simulation with 1000 realisations of MRW's of 4000 time steps with parameters extracted from the empirical data. Thus our proxy of multifractality appears to defy the hypothesis of a constant multifractal behaviour, as in the setup of the MRW model. The analysis has been carried out on a large set of data from different sectors of the market. A large number of stocks show patterns similar to those reported in figure 2.

We ascribe the inadequacy of the MRW to capture these observed empirical facts to the choice of a constant value of λ. As we have already mentioned above, a time varying $H(q)$ requires, through equation (6), a time varying λ. It seems in fact reasonable enough to suppose the scaling properties of financial time series to vary over time because of the reflection of complex and varying economic constraints.

We must remark that the values obtained for the correlation lengths T are very large compared to the time series lengths, a feature which is nonetheless commonly observed in multi-timescale volatility models [15]. The observed long-range nature of correlations seems to call for the introduction of non-stationary models, which of course would present a number of further inconveniences. However, multifractality is expected to be proportional only to λ^2 and thus the values of $\Delta H_{1,2}^w$ we measured should not be affected by the true T of the series. In fact, by taking equation (6) for two different values of q we have:

$$\zeta_{q_1}/q_1 - \zeta_{q_2}/q_2 = H(q_1) - H(q_2) \tag{13}$$

which gives after some rearrangement

$$\Delta H_{1,2} = \frac{\lambda^2}{2}(q_2 - q_1). \tag{14}$$

Therefore we expect our proxy of multifractality to increase along with an increase of λ, regardless of the measured T.

Note also that the limit of weak covariance, i.e. for $\lambda \to 0$, corresponds to a process whose behaviour is approximately unifractal. As one can appreciate from the quantiles reported in table III, the larger is λ, the larger are the GHE quantiles either, which reflect the irregularities present in the data. Note though that there is no correlation between the number of quantile-exceeding GHE's and λ. This fact tells us that a simple increase in λ would not necessarily make empirical data fit the model. It seems more reasonable instead to suppose the covariance structure of the log-volatility not to be constant over time, reflecting some underlying market mechanism.

A. The effect of t-Student residuals

Since the hypothesis of the residuals ϵ_k in equation (3) being Gaussian is at odds with the observed fat tails of the stock returns distributions, we have carried out



FIG. 2. *The black thick line represents the dynamical evolution of $\Delta H^w(1,2)$ ($\Delta H_t^w(1,2)$) for four stocks from different sectors of the market: Microsoft Corporation, IBM CO, Boeing Corporation and JP Morgan. $\Delta H_t^w(1,2)$ is evaluated on moving overlapping time windows of 1250 days, with a shift of 100 days. The first time window includes data from 1995 to 2000 and the corresponding $\Delta H_t^w(1,2)$ is plotted at the end of the period. The horizontal lines represent the 2.5% (red dashed line), 50% (green dot-dashed line) and 97.5% (magenta continuous line) quantiles extracted from the distribution of GHE estimated from 1000 realisations of MRW with parameters extracted from the data via the method described in the text. The quantile exceedances take place in all four cases with $\Delta H_t^w(1,2)$ always increasing on average. This suggests that in the period under study, the volatility may have exhibited a more complex behaviour than that predicted by constant multifractality models.*

the same analysis by taking the residuals to be t-Student distributed, that is

$$P(\epsilon) = \frac{1}{\sqrt{\pi}} \frac{\Gamma\left(\frac{1+\nu}{2}\right)}{\Gamma\left(\frac{\nu}{2}\right)} \frac{a^{\nu}}{(\epsilon^2 + a^2)^{\frac{1+\nu}{2}}}, \tag{15}$$

where ν is the number of degrees of freedom and the parameter a is related to the variance σ^2 via $\sigma^2 = a^2/(\nu-2)$. The student distribution is known to better fit stock returns tails, if one considers $\nu = 4$ or 5 [16]. The results obtained with this specification of the MRW model are reported in figure 4. We still retrieve a significant fraction of the observations overpassing the extreme quantiles, although this is reduced with respect to the case in which the residuals are Gaussian. We also observe a systematic shift upwards of the sample distributions of the $\Delta H^w(1,2)$'s, with the median now around 0.03. This tells us that, for a fixed covariance structure, the thickness of the tails dramatically rebounds on the multifractal properties of the simulated series, which have now a higher degree of multifractality, a feature which had also been reported in [6].

IV. DYNAMICAL WGHE ON MRW TIME SERIES

As a necessary test for the empirical analysis performed in the previous section, we now address the issue of studying the dynamical wGHE on synthetic time series generated via MRW. This model envisages a con-



FIG. 3. *We show how the quantiles of the wGHE distribution are affected by varying the correlation length T. We note no particular dependence, which in turn makes the test for varying multifractality depend only on λ.*

stant value for the scaling exponents and multifractality for given intermittency parameter λ and for each value of q, which means that possible variations of multifractality parameters for MRW times series are to be ascribed only to statistical fluctuations and are thus supposed not to cross the confidence intervals retrieved from their distributions. For this purpose, we test the wGHE method on a large set of MRW series to verify that the scaling exponents retrieved from these synthetic data are in agreement with the hypothesis of constant multifractality, which corresponds to fixing the set of parameters of the model $\{\lambda, T, \sigma\}$. In figure 5 we give examples of dynamical evolution $H^w(1)$ and $\Delta H^w_{1,2}$ that one observes for MRW series.

A complete list of the quantiles obtained for different values of λ is given in Table IV, together with the percentage of $\Delta H^w_{1,2}$ exceeding the confidence intervals. As also mentioned in the previous section, we have verified that the choice of T doesn't affect the quantile estimation: this is illustrated in figure 3, where we acknowledge that varying T, with λ fixed, doesn't affect the distribution of $\Delta H^w_{1,2}$. For this reason we fix $T = 500$ and vary λ. As one would expect, variations of these quantities occurring over time are well within the confidence intervals expected from the model. A comparison between the third column of table IV and the last right column of table III shows that, across different values of λ, the number of $\Delta^w_{1,2}$ exceeding the quantiles is much larger for empirical data than it is for synthetic MRW series. This analysis serves as a confirmation that the wGHE method returns values of multifractality and scaling exponents which are consistent with the hypothesis of the MRW and thus confirms further the anomalous nature of the fluctuations observed on empirical data.

V. CONCLUSIVE DISCUSSION

The scope of this paper has been to measure and validate variations of multifractality on a set of stock returns. As a benchmark we have considered the MRW, a parsimonious multifractal model that is able to reproduce faithfully many features commonly observed in stock returns. Since the dynamical estimations of multifractality are made on relatively small samples, to validate the observed fluctuations in the degree of multifractality as truly significant, we have first made an extensive study of the distribution of the generalised Hurst exponents $H(q)$ for different values of q. We have shown that the fluctuations observed in empirical data are truly crossing the extreme quantiles and we have checked that the GHE of MRW series fluctuates instead well within the confidence intervals expected for a multifractal process with constant degree of multifractality. The confidence intervals have been computed on many realisations of MRW's simulated with a set of parameters obtained from different empirical data series. We ascribe the disagreement between model and data to the intermittency parameter being constant. We have shown that increasing the intermittency parameters makes the tails of the GHE distribution computed on the synthetic series thicker, while we found no relevant dependence with T. Our analysis thus suggests that a varying intermittency parameter may be the correct guess towards the inclusion of the observed empirical facts into the MRW model. A possible extension, which we are investigating at the present, is the possibility of identifying switching points between multifractality regimes, a topic which could be steered along the research stream of regime switching in dynamical correlation [26, 27]. The idea is to look for a covariance structure of the form

$$\begin{aligned}\mathrm{Cov}(\omega_k, \omega_{k+h}) =& \Theta(k-k^*)\lambda_1^2 \log \frac{T_1}{(1+|h|)\Delta t} \\ &+ \Theta(k^*-k)\lambda_2^2 \log \frac{T_2}{(1+|h|)\Delta t}, \qquad (16)\end{aligned}$$

where $\Theta(x)$ denotes the Heaviside function. The subtlety of this approach is nevertheless the inclusion of a new parameter k^* defining the switching point between different covariance regimes which, although stimulating for its possible economic meaning, seems to be tricky to be measured.

The finding of very large correlation lengths is, in this framework, probably the best clue one has to detect nonstationarity in financial time series. The very large memory present in the volatility is an intriguing concept which could also be related to the way markets evolves with history: the hypothesis of having a varying multi scaling behaviour might suggest that memory is itself a time varying process whose prolongation is to be ascribed to the growing complexity of modern societies which financial markets reflect. In this regard, a more detailed study of the spectrum of the covariance structure of the volatility on very large time series is likely to reveal some new



FIG. 4. *The dynamical evolution of $\Delta H^w(1,2)$ for the four companies IBM Corp, American Express, Boeing Corp and Bank of America. Now the quantiles have been obtained by considering a MRW with t-Student residuals. The distributions of the simulated returns have thicker tails than in the original Gaussian case, and this results in the quantiles of the sample distributions of $\Delta H^w(1,2)$ being much larger. Nonetheless, the fluctuations observed on empirical data are still overpassing the quantiles significantly: 25% (IBM), 21% (Bank of America), 14% (American Express) and 18% (Boeing). We also note a significant shift upwards of the median of the sample distributions of $\Delta H^w(1,2)$. This is a clue that multifractality is reflecting the tails of the synthetic returns distribution, although the covariance structure stays the same.*

insights (see [28] for a relevant discussion on this matter). The GHE method has been proved to be the most efficient among other scaling measures [4]; nonetheless, it would be interesting to investigate whether it is possible to retrieve convincing evidence of variations in multifractality by means of other methods. The study of the unconditional distribution of returns at all time scales could also be performed by means of parametric methods; these have the drawback of requiring a specific model to be chosen though and, given the non-stationary behaviour of financial time series, may be inappropriate, apart from very few practical situations. When one needs to carry out dynamic maximum likelihood estimation on the empirical data over the time windows at different time scales, the shortness of the time windows, especially at large time scales, results in the estimator not to converge and in the results to be hardly interpretable. Starting from the results of this work, our main future aim is the development of a realistic model for stock dynamics which incorporate these empirical findings, possibly backed by a plausible economic ground about what triggers multifractality to change over time.

Appendix A: Generalized Hurst exponent method

The generalized Hurst exponent $H(q)$ is defined when the following scaling is observed:

$$K_q(\tau) \propto \tau^{qH(q)} \quad , \tag{A1}$$



FIG. 5. *Left: Dynamical evolution (black thick line) of the wGHE* $H^w(q=1)$ *computed on a realisation of a synthetic MRW time series. The horizontal lines represent the 2.5% (red dashed line), 50% (green dot-dashed line) and 97.5% (magenta continuous line) quantiles extracted from the distribution of GHE estimated from 1000 realisations of MRW with parameters* $\lambda = 0.2$*, T=500,* $\sigma = 1$*. Although statistical fluctuations are present they stay within the confidence intervals corresponding to the chosen set of parameters. Right: Dynamical evolution of* $\Delta H^w_{1,2}$ *on another realisation of MRW with same parameters as in the left panel. As for the scaling exponent* $H^w(q=1)$*, the fluctuations are compatible with a constant multifractality scenario. In both panels the shaded area represents the standard error respectively on the wGHE and on* $\Delta H^w_{1,2}$*. The latter is computed as* $\boldsymbol{std}\left(\Delta H^w_{1,2}\right) = \sqrt{var(H^w(1)) + var(H^w(2))}$*, assuming that* $\boldsymbol{cov}(H^w(1), H^w(2)) = 0$*.*

TABLE IV. *Quantiles and number of* $\Delta H^w_{1,2}$ *exceedances obtained from several simulations of MRW with different* λ*. Note that the percentage of* $\Delta H^w_{1,2}$ *exceeding the quantiles is negligible. This proves that the dynamic wGHE is faithfully reproducing the constant multifractality envisaged by the model. In all simulations we fixed* $T = 500$ *and* $\sigma = 1$*, after verifying that this choice does not affect the results: in fact, all different values of* T *and* σ *which we have tried return the same kind of percentages of exceedance.*

λ	Quantiles	Exceedances percentage
0.12	{-0.0234,0.0033,0.0253}	0 %
0.14	{-0.0227,0.0045,0.0258}	1 %
0.16	{-0.0221,0.0048,0.0276}	1 %
0.18	{-0.0223,0.0052,0.0272}	1 %
0.20	{-0.0203,0.0056,0.0302}	0 %
0.22	{-0.0244,0.0066,0.0310}	0 %
0.24	{-0.0201,0.0076,0.0314}	5 %
0.26	{-0.0211,0.0078,0.0330}	1 %
0.28	{-0.0238,0.0081,0.0334}	0 %
0.30	{-0.0219,0.0096,0.0357}	1 %

where $K_q(\tau)$ are the qth-order moment of the increment of a process $X(t)$, defined as [29]

$$K_q(\tau) = \frac{\langle |X(t+\tau) - X(t)|^q \rangle}{\langle |X(t)|^q \rangle}. \tag{A2}$$

τ can vary between 1 and τ_{max} and $\langle\cdot\rangle$ denotes the sample average over the time-window. Equation (A1) provides a way to compute directly the GHE for any value of q, as long as the scaling is observed.

Financial returns time series exhibit often very large fluctuations, corresponding to periods of abnormal activity or large scale financial crises. These fluctuations may severely flaw any statistical estimator, when the latter is computed dynamically and the number of observations is therefore reduced; to track the dynamical evolution of the GHE, an exponential smoothing for the moments has been introduced in [23], with corresponding weights chosen as the best trade-off between length of the time window and sensitivity to possible changes in scaling properties of the time series. Details on the optimal weights can be found in [23, 30]. Equation (A2) is then replaced by

$$K_q^w(t,\tau) = \frac{\langle |S(t+\tau) - S(t)|^q \rangle_w(t)}{\langle |S(t)|^q \rangle_w(t)}, \qquad \text{(A3)}$$

where $\langle\cdot\rangle_w$ denotes weighted average. Analogously the weighted GHE (wGHE) $H^w(q)$ is computed by fitting the linear relation obtained inverting the scaling relation observed for the weighted moments:

$$K_q^w(\tau) \propto \tau^{qH^w(q)}. \qquad \text{(A4)}$$

The exponent $H^w(q)$ embodies the scaling behaviour of the corresponding q-moment when the time series is exponentially smoothed.

Appendix B: Validity of the scaling

A more detailed comparison between the relations (7) and the the GHE's obtained from simulated series with different values of λ has also been performed. We have simulate many synthetic MRW time series with λ varying in $[0, 1]$ and for each we have computed the GHE. In figure (6) we plot the relations we obtain for $H(q = 1, 3)\, vs\, \lambda$. The black solid line in both figures is the best quadratic fit obtained from the measured points. We retrieve, as expected from equations (7), quadratic trends for both $H(1)$ and $H(3)$ *vs* λ, although the actual slope is not consistent with that predicted theoretically. We find in fact

$$H(1) = \frac{1}{2}(1 + a\lambda^2), \quad H(3) = \frac{1}{2}(1 - b\lambda^2) \qquad \text{(B1)}$$

with $a = 0.1348$ and $b = 0.2374$ in the figure. Repeating the experiment many times we obtain $a = 0.1475 \pm 0.011$ and $b = 0.2502 \pm 0.017$, still very far from the theoretical values $a = b = 1$. This apparent disagreement between the GHE method and the theoretical scaling function is probably due to the growing amount of noise present in the synthetic series as one increases λ. Better fits are retrieved in fact if we shrink the upper bound of λ: fits restricted to values of $\lambda < 0.3$ work already much better giving $a = 0.7162 \pm 0.012$ and $b = 0.5512 \pm 0.018$. Note also that financial data exhibit, most of the time, intermittency parameters $\lambda < 0.3$ (see also [31]). Somewhere around $\lambda \sim 0.25$ though, both fits start not to be distinguishable from a flat trend at $H(1) = 1/2$ and $H(3) = 1/2$, as expected for $H(2)$. We conclude that the noise present in the GHE doesn't allow to estimate the intermittency parameter reliably from the structure functions.

Nonetheless this apparent flaw of the method doesn't affect the validity of the empirical results presented in section III, since we chose to estimate the intermittency parameter from the volatility autocorrelation rather than from the scaling functions, although both methods being in principle valid.

ACKNOWLEDGMENTS

The authors warmly thank Jean-Philippe Bouchaud for helpful discussions.

---

[1] J. Kantelhardt, S. Zschiegner, E. Koscielny-Bunde, S. Havlin, A. Bunde, and H. Stanley, Physica A **316**, 87 (2002).

[2] T. Matteo, T. Aste, and M. Dacorogna, Journal of Banking & Finance **29**, 827 (2005).

[3] J. Muzy, J. Delour, and E. Bacry, The European Physical Journal B **17**, 537 (2000).

[4] J. Barunik and L. Kristoufek, Physica A **389**, 3844 (2010).

[5] T. Di Matteo, Quantitative Finance **7**, 21 (2007).

[6] J. Barunik, T. Aste, T. Di Matteo, and R. Liu, Physica A **391**, 4234 (2012).

[7] L. Kristoufek, Europhysics Letters **95**, 68001 (2011).

[8] E. Bacry, J. Delour, and J. Muzy, Physical Review E **64**, 026103 (2001).

[9] J. Bouchaud, M. Potters, and M. Meyer, The European Physical Journal B **13**, 595 (2000).

[10] R. Liu, T. Matteo, and T. Lux, Advances in Complex Systems **11**, 669 (2008).

[11] R. Liu, T. Di Matteo, and T. Lux, Physica A **383**, 35 (2007).

[12] B. Mandelbrot, A. Fisher, and L. Calvet, Cowles Foundation discussion paper no. 1164, Yale University (1997).

[13] T. Lux, Journal of business & economic statistics **26**, 194 (2008).

[14] Z. Ding, C. Granger, and R. Engle, Journal of empirical finance **1**, 83 (1993).

[15] L. Borland and J. Bouchaud, Science & Finance (CFM) working paper archive (2005).

[16] J. Bouchaud and M. Potters, *Theory of financial risk and derivative pricing: from statistical physics to risk management* (Cambridge Univ Pr, 2003).

[17] T. Di Matteo, T. Aste, and M. Dacorogna, Physica A **324**, 183 (2003).

[18] M. Bartolozzi, C. Mellen, T. Di Matteo, and T. Aste, The European Physical Journal B **58**, 207 (2007).

[19] W. Zhou, EPL **88**, 28004 (2009).

[20] M. Dacorogna, *An introduction to high-frequency finance* (Academic Pr, 2001).

[21] A. Carbone, G. Castelli, and H. Stanley, Physica A **344**, 267 (2004).

[22] D. Grech and G. Pamuła, Physica A **387**, 4299 (2008).



FIG. 6. *We plot the $H(q)$ vs λ for $q = 1$ (left) and $q = 3$ (right). The dots refer to the $H(q)$'s values obtained from the MRW-simulated time series with λ ranging in [0,1]. The black solid lines are the best fits to equations B1.*

[23] R. Morales, T. Di Matteo, R. Gramatica, and T. Aste, Physica A **391**, 3180 (2012).

[24] A. Arneodo, E. Bacry, S. Manneville, and J. Muzy, Physical Review Letters **80**, 708 (1998).

[25] L. Borland, J. Bouchaud, J. Muzy, and G. Zumbach, *The dynamics of financial marketsMandelbrots multifractal cascades*, Tech. Rep. (and beyond. Working Paper, 2008 (Science & Finance (CFM) working paper archive (500061)), 2005).

[26] D. Pelletier, Journal of Econometrics **131**, 445 (2006).

[27] J. Hamilton, Journal of Econometrics **45**, 39 (1990).

[28] R. Chicheportiche and J. Bouchaud, Journal of Statistical Mechanics **P09003** (2011).

[29] A.-L. Barabasi and T. Vicsek, Phisical Review A **44**, 2730 (1991).

[30] F. Pozzi, T. Di Matteo, and T. Aste, The European Physical Journal B **85**, 175 (2012).

[31] E. Bacry, A. Kozhemyak, and J. Muzy, Journal of Economic Dynamics and Control **32**, 156 (2008).